

DR INGRID DEMMELMAIER (Orcid ID : 0000-0002-2068-4708)

MRS ANN CHRISTIN HELGESEN BJØRKE (Orcid ID : 0000-0002-5238-9826)

DR LAURIEN M BUFFART (Orcid ID : 0000-0002-8095-436X)

Article type : Original Article

Does exercise intensity matter for fatigue during (neo-)adjuvant cancer treatment? The Phys-Can randomised clinical trial

Ingrid Demmelmaier, Hannah L Brooke, Anna Henriksson, Anne-Sophie Mazzoni, Ann Christin Helgesen Bjørke, Helena Igelström, Anna-Karin Ax, Katarina Sjövall, Maria Hellbom, Ronnie Pingel, Henrik Lindman, Silvia Johansson, Galina Velikova, Truls Raastad, Laurien M Buffart, Pernilla Åsenlöf, Neil K Aaronson, Bengt Glimelius, Peter Nygren, Birgitta Johansson, Sussanne Börjeson, Sveinung Berntsen, Karin Nordin

Note: Ingrid Demmelmaier and Hannah L Brooke should be considered joint first author

Author institutions and affiliations

PhD Ingrid Demmelmaier, PhD Hannah L Brooke, PhD Anna Henriksson, MSc Anne-Sophie Mazzoni, Professor Karin Nordin, Uppsala University, Department of Public Health and Caring Sciences, Uppsala, Sweden

MSc Ann Christin Helgesen Bjørke, Professor Sveinung Berntsen, University of Agder, Department of Sport Science and Physical Education, Kristiansand, Norway

PhD Helena Igelström, Professor Pernilla Åsenlöf, Uppsala University, Department of Neuroscience, Uppsala, Sweden

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/SMS.13930](https://doi.org/10.1111/SMS.13930)

This article is protected by copyright. All rights reserved

MSc Anna-Karin Ax, Professor Sussanne Börjeson, Linköping University, Department of Oncology and Department of Medical and Health Sciences, Linköping, Sweden

PhD Katarina Sjövall, Lund University, Department of Oncology and Skåne University Hospital, Department of Oncology, Lund, Sweden

PhD Maria Hellbom, Stockholm Health Care Services, Centre for Cancer Rehabilitation, Stockholm, Sweden

PhD Ronnie Pingel, Uppsala University, Department of Statistics, Uppsala, Sweden

PhD Henrik Lindman, PhD Silvia Johansson, Professor Bengt Glimelius, Professor Peter Nygren, PhD Birgitta Johansson, Uppsala University, Department of Immunology, Genetics and Pathology, Uppsala, Sweden

Professor Galina Velikova, University of Leeds, Leeds Cancer Centre, Leeds Institute of Medical Research at St James's, Leeds, UK

Professor Truls Raastad, Norwegian School of Sport Sciences, Department of Physical Performance, Oslo and University of Agder, Department of Sport Science and Physical Education, Kristiansand, Norway

PhD Laurien M Buffart, Radboudumc, Department of Physiology, Nijmegen, The Netherlands

Professor Neil K Aaronson, The Netherlands Cancer Institute, Division of Psychosocial Research & Epidemiology, Amsterdam, The Netherlands

Corresponding author

PhD Ingrid Demmelmaier, Uppsala University, Department of Public Health and Caring Sciences, Husargatan 3, Post Box 564, 751 22 Uppsala, Sweden

ingrid.demmelmaier@pubcare.uu.se

Acknowledgements

The authors would like to acknowledge the study participants, the clinicians involved in recruitment, the staff at the public gyms where the resistance exercise was performed, and the patient representatives who contributed with their perspectives. We also

Accepted Article

acknowledge Dr Pernille Højman, Centre for Physical Activity Research and Centre of Inflammation and Metabolism, Rigshospitalet Copenhagen, Denmark, who contributed substantially to the project but sadly passed away in April 2019.

This work was supported by The Swedish Cancer Society (grant numbers 150841, 160483); The Swedish Research Council (grant number KDB/9514); The Nordic Cancer Union (2015), and The Oncology Department Foundations Research Fund in Uppsala (2016, 2017).

The authors declare they have no conflict of interest.

List of abbreviations in text

CRF = Cancer-Related Fatigue

BCS = Behaviour Change Support

MFI = Multidimensional Fatigue Inventory

FACIT-F = Functional Assessment of Chronic Illness Therapy-Fatigue scale

HRQoL = Health-Related Quality-of-Life

BMI = Body Mass Index

VO₂max = Maximal Oxygen uptake

Borg RPE = Borg Rating of Perceived Exertion

RER = Respiratory Exchange Ratio

HADS = Hospital Anxiety and Depression Scale

EORTC QLQ-C30 = European Organisation for Research and Treatment of Cancer,
Quality of life Questionnaire C30

MVPA = Moderate to Vigorous intensity Physical Activity

ABSTRACT

Exercise during cancer treatment improves cancer-related fatigue (CRF), but the importance of exercise intensity for CRF is unclear. We compared the effects of high vs. low-to-moderate intensity exercise with or without additional behaviour change support (BCS) on CRF in patients undergoing (neo-)adjuvant cancer treatment. This was a multicentre, 2x2 factorial design randomised controlled trial (Clinical Trials NCT02473003) in Sweden. Participants recently diagnosed with breast (n=457), prostate (n=97) or colorectal (n=23) cancer undergoing (neo-)adjuvant treatment were randomised to high intensity (n=144), low-to-moderate intensity (n=144), high intensity with BCS (n=144) or low-to-moderate intensity with BCS (n=145). The 6-month exercise intervention included supervised resistance training and home-based endurance training. CRF was assessed by Multidimensional Fatigue Inventory (MFI, five subscales score range 4-20), and Functional Assessment of Chronic Illness Therapy-Fatigue scale (FACIT-F, score range 0-52). Multiple linear regression for main factorial effects was performed according to intention-to-treat, with post-intervention CRF as primary endpoint. Overall, 577 participants (mean age 58.7 years) were randomised. Participants randomised to high vs. low-to-moderate intensity exercise had lower physical fatigue (MFI Physical Fatigue subscale; mean difference -1.05 [95% CI:-1.85,-0.25]), but the difference was not clinically important (i.e.<2). We found no differences in other CRF dimensions and no effect of additional BCS. There were few minor adverse events. For CRF, patients undergoing (neo-)adjuvant treatment for breast, prostate or colorectal cancer can safely exercise at high or low-to-moderate intensity, according to their own preferences. Additional BCS does not provide extra benefit for CRF in supervised, well-controlled exercise interventions.

Keywords

Endurance training, resistance training, oncology, behaviour change, cancer-related fatigue.

INTRODUCTION

Cancer survival rates have improved due to earlier detection and advances in treatment.¹ However, cancer survivors report long-term challenges, such as cancer-related fatigue (CRF), physical deconditioning, and decreased health-related quality-of-life (HRQoL).² CRF is defined as a distressing, persistent sense of physical, emotional and/or cognitive tiredness or exhaustion that is not proportional to recent activity and interferes with usual functioning.³ Prevalence of moderate-to-severe CRF during treatment is 30-60%,³ and clinically important CRF has been reported in one-third of patients up to 6 years after treatment.⁴ Symptom clusters of co-occurring CRF, pain and psychological distress may be persistent⁵ and are associated with lower HRQoL.⁶ The aetiology of CRF is multifactorial and treatment-induced activation of pro-inflammatory cytokines may be one trigger.³

Exercise during and after treatment is effective in counteracting CRF,⁷ possibly by lowering the inflammatory activity and/or by increasing physical fitness.⁸ Exercise also improves HRQoL,⁹ may increase chemotherapy completion rates,^{10,11} and reduce the risk of cancer mortality.¹² International guidelines¹³ recommend 3 sessions of at least moderate intensity endurance training and/or 2 sessions of at least moderate intensity resistance training each week to counteract CRF. However, the evidence-base regarding the ideal 'exercise prescription' in terms of exercise frequency, intensity, duration, and type for cancer survivors is insufficient.¹³ While low-to-moderate intensity may be preferred by most patients,¹⁴ one study found that moderate-to-high intensity exercise during cancer treatment was beneficial for physical fatigue compared to low intensity exercise.¹⁰ However, exercise volume was not controlled for, so the importance of intensity *per se* could not be determined. Second-generation studies comparing different exercise intensities in relation to side-effects, such as CRF, are therefore needed.¹³

Many patients find it difficult to perform physical activity during oncological treatment and they are on average less physically active than the general population.¹⁵ In exercise interventions, adherence rates have been reported between 23%¹⁶ and 84%,¹⁷ depending on the characteristics of the sample, the intervention and the method used to calculate adherence. Interview studies indicate a need for individualised support to overcome barriers such as side-effects and external demands.¹⁸ Behaviour change

support (BCS), can be used to overcome such barriers. Systematic reviews and meta-analysis have identified behaviour change strategies associated with high adherence and larger effect sizes in interventions designed to increase physical activity among cancer survivors. Examples of such techniques are self-monitoring,¹⁹ goal-setting,²⁰ graded tasks,^{20,21} social support²² and supervision.^{20,22} However, if BCS can influence health outcomes through increased intervention adherence is unclear due to limitations in study methodology and reporting.^{20,23}

The primary aim of this second-generation study was to determine the effects of high vs. low-to-moderate intensity exercise with or without additional BCS on CRF (primary endpoint) in patients undergoing (neo-)adjuvant cancer treatment. Secondary aims were to determine the effects on HRQoL, anxiety/depression, function in daily life, cardiorespiratory fitness, muscle strength, level of physical activity, sedentary time, sleep and treatment completion rates.

METHODS

Design

The Physical training and Cancer (Phys-Can) study was a Swedish three-centre, 2x2 factorial design randomised controlled trial (Figure 1), previously described in depth.²⁴ Briefly, participants were randomised to high or low-to-moderate intensity exercise, with or without additional BCS (ClinicalTrials.gov NCT02473003). The Swedish Ethical Review Authority approved the study (Dnr 2014/249).

Coaches (qualified and experienced physiotherapists, n=13 or personal trainers, n=2) were assigned to lead an intervention group based on logistics, scheduling and their additional competence in BCS. Each coach supervised participants in both exercise intensity groups. However, coaches who provided additional BCS supervised only those groups.

There were no major changes to the study after the trial commenced.

Participants

Participants were recruited from Uppsala, Lund and Linköping University hospitals from March 2015 to April 2018. Eligible participants, assessed by an oncologist, were >18 years, literate in Swedish and recently diagnosed with curable breast (women only), prostate or colorectal cancer, scheduled to begin (neo-)adjuvant chemotherapy, radiotherapy and/or endocrine therapy. The chemotherapy treatment period was typically 4 months for breast cancer and 6 months for colorectal cancer. Radiotherapy treatment was typically 2 months for prostate cancer and 3 weeks for breast cancer. Endocrine therapy were scheduled for 2.5-10 years and antibody treatment for 6-12 months.

Exclusion criteria were stage IIIb-IV breast cancer, inability to perform basic activities of daily living, cognitive disorders, severe psychiatric disease or other disabling conditions that might contraindicate high intensity exercise (e.g. severe heart failure, severe chronic obstructive pulmonary disease or orthopaedic conditions), treatment for an additional ongoing malignant disease, BMI<18.5 kg/m² or pregnancy. A research nurse/assistant provided oral and written information to eligible participants prior to start of treatment. Those willing to participate gave written informed consent before baseline data collection.

Interventions

The 6-month intervention was initiated at start of the (neo-)adjuvant cancer treatment as described in Table 1. The supervised, group-based resistance training at public gyms was performed twice/week (Suppl. Figure 1). The home-based endurance training at high intensity consisted of interval training, performed twice/week, while low-to-moderate intensity endurance training consisted of 150 weekly minutes of walking or biking. Additional BCS, such as goal-setting, planning and self-monitoring, was delivered face-to-face jointly with the resistance training sessions. Patients with breast cancer, scheduled for neo-adjuvant chemotherapy, exercised during the four months pre-surgery only. Standardised delivery of both exercise and additional BCS was assessed and enhanced as described in Table 1.

The intervention was developed by researchers with expertise in exercise physiology, BCS and physiotherapy in collaboration with clinicians (oncologists and physiotherapists) and patient representatives.

Outcomes and data management

Follow-up data collection was completed in November 2018. CRF was assessed with the Multidimensional Fatigue Inventory (MFI),²⁵ measuring General, Physical and Mental Fatigue, Reduced Motivation and Reduced Activity, each subscale range 4-20. Based on previous research,¹⁰ the Physical Fatigue subscale was used for power calculation.

CRF was also assessed with Functional Assessment of Chronic Illness Therapy – Fatigue (FACIT-F) scale,²⁶ a frequently used single scale questionnaire with range 0-52.

All outcomes were assessed with well-established and validated methods completed at home, at baseline (before randomisation) and immediately post-intervention. HRQoL was assessed with European Organisation for Research and Treatment of Cancer (EORTC) QLQ-C30,²⁷ anxiety and depression with Hospital Anxiety and Depression Scale (HADS)²⁸ and functioning in daily life with World Health Organization Disability Assessment Schedule (WHODAS) 2.0 Work and Social Participation subscales.²⁹ Muscle strength was assessed with 1 repetition maximum (RM) test of upper and lower extremities. Chemotherapy completion rate was reported as relative dose intensity based on data from medical records.³⁰ For details about assessment methods, see Suppl. Table 1. Composite scores for questionnaires were calculated according to published instructions.

Physical activity was measured using SenseWear Armband mini (BodyMedia Inc., Pittsburgh, PA, USA) at baseline and post-intervention, as described in Suppl. Table 1. A minimum of 80% wear time for four out of seven days was required for data to be included. Mean sedentary time per 24 hrs was considered to be time spent at 0-1.5 METs according to SenseWear algorithms and mean time in moderate to vigorous intensity physical activity (MPVA) per 24 hrs was considered to be that of at least 3 METs according to SenseWear algorithms.³¹

Cardiorespiratory fitness was independently assessed in dedicated exercise labs using maximal oxygen uptake (VO_{2max}) tests with gas exchange measurement, walking/running to exhaustion using a modified Balke protocol.³² Tests were accepted if two out of three criteria were fulfilled: 1) tester judged the test as maximal, 2) Borg RPE³³ rating ≥ 17 and 3) respiratory exchange ratio (RER) ≥ 1.1 .³⁴

Exercise adherence was recorded as absolute numbers of performed sessions and calculated as % volume. Exercise volume (frequency, intensity and time) was calculated as performed exercise divided by prescribed (maximum possible) exercise, in accordance with previous research.³⁵ This calculation resulted in one proportion (0-100%) for resistance training and one proportion (0-100%) for endurance training for each participant. These proportions were pooled to mean adherence per exercise component and intervention group. For details, see Table 1 under the heading "Calculation of adherence". **Sample size**

An *a-priori* power calculation showed that 600 participants (150 per trial arm) were required to detect main factorial effects of exercise intensity and BCS on the MFI Physical Fatigue subscale,³⁶ with a minimum clinically important difference of 2 points (SD 5) post-intervention and 80% power at alpha level 0.05. This calculation allowed for missing data and drop-out from the study.²⁴

Randomisation

The random allocation sequence with a ratio of 1:1:1:1 was computer generated and thus concealed from all research staff. Within each stratum (3 centres and 3 diagnoses) randomisation was carried out following a permuted block design with 8 participants per block. Once baseline data collection was finalised, each participant was automatically assigned to an intervention group using a web-portal.

Blinding

Blinding of coaches and participants to the intervention group was not feasible. However, coaches and participants were informed that there was limited evidence for which intensity would be more beneficial for CRF.

Statistical methods

Differences in baseline characteristics between those included in analysis and those who withdrew were examined using t-tests for continuous variables and chi-squared tests for categorical variables.

Multiple linear regression was used to simultaneously estimate the main effect for exercise intensity (high vs. low-to-moderate intensity) and BCS (with vs. without), and their interaction (intensity x BCS) on each outcome post-intervention. Results are presented as adjusted mean difference with 95% Confidence Intervals (95%CI). Analyses were conducted according to intention-to-treat. Models included baseline measures of the outcome to increase precision and were adjusted for centre and cancer diagnosis.

Missing data were accounted for using multiple imputations by chained equations. Auxiliary variables used to inform imputed values were age, education level, centre, diagnosis, chemotherapy treatment, baseline values of outcome measure and intervention group. Where missing data at baseline were >10%, baseline data for that outcome were not included as an auxiliary variable or as a variable in the main models, as baseline data was deemed not to add additional precision for these outcomes.^{37,38}

Differences between intervention groups in adherence to resistance and endurance training volume were examined using one-way ANOVA. Among participants in the low-to-moderate intensity groups minutes of exercise above the prescribed intensity (i.e. >60% of heart rate reserve) were compared between those who did and did not receive additional BCS using a t-test.

A supplementary analysis based on complete cases was performed. In addition, each cancer diagnosis was examined separately; these models were adjusted for centre and baseline measures of the outcome variable.

Analyses were carried out in Stata version 15.0.

Patient and public involvement

Patient representatives, one from each diagnosis group, were included in the project group. They were involved in the design of the study, the content of the intervention, informational material, and provided feedback on the burden of the intervention. They will help disseminate the results within their respective patient organisation.

RESULTS

Recruitment

Six hundred (29%) of 2051 eligible patients agreed to participate (Figure 1). Participation rate per diagnosis was 30% for breast cancer, 22% for prostate cancer and 20% for colorectal cancer. Participants, compared to those who declined participation, were younger (mean age 58.7 vs 63.6 years, p -value<0.001) and less likely to have prostate or colorectal cancer than breast cancer (OR [95%CI]; colorectal cancer, 0.58 [0.36-0.92]; prostate cancer, 0.65 [0.51-0.83]). Twenty-three participants withdrew from the study before randomisation. In total, 577 participants were randomised.

Baseline characteristics of randomised participants

Breast cancer was the most common diagnosis ($n=457$), followed by prostate cancer ($n=97$) and colorectal cancer ($n=23$). Most participants with breast (84.5%) or prostate (66.7%) cancer received combinations of (neo-)adjuvant treatments. Among these participants, the most common combinations were chemotherapy with radiotherapy and endocrine treatment for breast cancer (30.4%) and combined neo-adjuvant and adjuvant endocrine treatment for prostate cancer (51.5%). All participants with prostate cancer also received curative radiotherapy. All participants with colorectal cancer received adjuvant chemotherapy (Table 2). Time between diagnosis and randomisation was median 61 (interquartile range 44-86) days and there was a median of 7 (interquartile range 5-11) days between randomisation and starting the intervention. Sociodemographic, disease and treatment characteristics, as well as physical fitness and physical activity levels, were similar across all four intervention groups (Table 2 and Suppl. Table 2).

Missing data

Overall, 89 randomised participants withdrew from the study before the follow-up assessment. The reasons for withdrawal were too far to travel/lack of time ($n=36$), illness/side-effects of treatment ($n=25$) and lack of motivation/disliked training ($n=10$), with similar distribution between intervention groups. Eighteen participants did not

provide a reason. Missing baseline data on outcomes ranged from n=20 (FACIT-F) to n=113 (VO₂max) (Suppl. Table 3). Missing data were not associated with intervention group, age, living situation, education, weight status, and general, physical or mental fatigue. However, individuals who were not included in the analysis of the main outcome had slightly lower cardiorespiratory fitness (VO₂max mean [SD]: 29.2 [7.5] vs 30.8 [7.0] ml/kg/min p-value=0.043) and HRQoL (EORTC QLQ-C30 summary score mean [SD]: 80.6 [13.7] vs 83.7 [12.0] p-value=0.013) at baseline compared to those included.

Primary outcome

Participants randomised to exercise at high compared with low-to-moderate intensity had lower MFI physical fatigue (adjusted mean difference -1.05 [95% CI, -1.85 to -0.25]) (Table 4), while there were no differences for the other MFI subscales or for FACIT-F. Moreover, there were no main effects for additional BCS and no exercise intensity-BCS interactions for any CRF measures.

Secondary outcomes

There were no main effects of exercise intensity or additional BCS on HRQoL based on the EORTC QLQ C-30 Summary Score (Table 4). However, there was an interaction effect indicating that in the groups receiving additional BCS, high intensity exercise was associated with a lower HRQoL compared to low-to-moderate intensity, while in the groups not receiving BCS high intensity exercise was associated with higher HRQoL compared to low-to-moderate intensity (Table 4 and Suppl. Table 4).

Participants randomised to exercise at high vs. low-to-moderate intensity had better cardiorespiratory fitness (adjusted mean difference 1.61 [95% CI 0.19-3.04] ml/kg/min) and greater leg strength (adjusted mean difference 3.98 [95% CI 0.58-7.38] kg) (Table 4). There were no main or interaction effects of additional BCS for these outcomes. An exercise intensity-BCS interaction was observed for MVPA (Table 4).

There were no main or interaction effects of the intervention on anxiety, depression, functioning in daily life, sleep, sedentary behaviour, chemotherapy completion rates or relative dose intensity (Table 4).

For chemotherapy completion rates, 118 participants (39.6%) out of 298 had dose reduction or treatment discontinuation. Mean relative dose intensity was 90.2% in high intensity with BCS, 92.6% in high intensity without BCS, 93.6% in low-to-moderate intensity with BCS, and 92.9% in low-to-moderate intensity without BCS (Suppl. Table 5). There were no adjustments of radiotherapy dose for any participant.

Supplementary analyses

The results from the complete case analysis were similar to the main results, i.e. the MFI Physical Fatigue scale differed significantly, in favour of the high intensity exercise

groups (Suppl. Table 3). However, in contrast to the intention-to-treat analysis, VO_{2max} did not differ between groups in the complete cases analysis. Diagnosis-specific analyses reflected the main results for all diagnostic groups (Suppl. Tables 6 and 7) but were underpowered for participants with colorectal cancer and weaker for those with prostate cancer.

Intervention adherence

Participants completed on average 50.4% of the prescribed resistance training volume, with no differences between intervention groups ($p=0.438$) (Figure 2). For attendance specifically, the absolute mean (SD) number of performed sessions during the intervention period was 25 (10). Reported per group, mean (SD) number of performed sessions was 23 (10) for high intensity with BCS, 24 (9) for high intensity without BCS, 26 (10) for low-to-moderate intensity with BCS, and 26 (8) for low-to-moderate intensity without BCS.

Adherence to home-based endurance training volume differed between groups (mean [SD] % were 38.8 (33.1) for high intensity with BCS, 41.6 (33.6) for high intensity without BCS, 57.7 (38.3) for low-to-moderate intensity with BCS, and 51.4 (38.7) for low-to-moderate intensity without BCS, $p<0.001$) and pair-wise comparisons are presented in Figure 2. The absolute mean (SD) number of performed interval sessions during the intervention period was 23 (17) for high intensity with BCS and 23 (16) for high intensity without BCS. The mean (SD) number of continuous training sessions (minimum 10-minute bouts), was 73 (66) for low-to-moderate intensity with BCS and 70 (61) for low-to-moderate intensity without BCS. Among participants in the low-to-moderate intensity groups those who received additional BCS performed more minutes of exercise above the prescribed intensity (i.e. $>60\%$ of heart rate reserve) than those who did not receive additional BCS (506 vs 326 minutes, $p\text{-value}=0.026$).

Adverse events due to intervention

Thirty-two minor adverse events in 30 participants ($n=8$ high intensity with BCS, $n=12$ high intensity without BCS, $n=6$ low-to-moderate with BCS, $n=4$ low-to-moderate without BCS) prevented them from completing the ongoing training session. These events

included muscle strains, joint pain, and dizziness. In addition, three participants needed to attend hospital as a result of exercise; one injured a finger, and two fainted.

Accepted Article

DISCUSSION

This was a large, second-generation RCT designed to examine the effects of exercise intensity *per se* on CRF and other health outcomes during cancer treatment. High intensity resistance and endurance exercise yielded significantly lower physical fatigue compared to low-to-moderate intensity exercise in patients undergoing (neo-)adjuvant treatment, but the magnitude of effect did not reach the minimal clinically important difference of two points.³⁶ Further, there were no differences between groups in other CRF dimensions. There were few minor adverse events, which indicates that exercise is safe, even at high intensity, for these patient groups. Although there were small benefits of high intensity exercise for muscle strength and cardiorespiratory fitness, overall, patients undergoing (neo-)adjuvant treatment for breast, prostate or colorectal cancer can be advised to exercise at either intensity, according to their own preferences. Finally, our results suggest that, in a motivated and relatively healthy sample, additional BCS does not influence CRF.

In line with our results, van Waart et al demonstrated statistically lower physical fatigue after combined resistance and endurance exercise at moderate-to-vigorous intensity compared to low intensity walking during adjuvant chemotherapy,¹⁰ but did not find a minimum clinically important difference between groups.³⁶ However, van Waart et al did not control for exercise volume, which limited the possibilities to draw conclusions about the effect of exercise intensity *per se*.

Kampshoff et al reported no between-group differences in CRF for high vs low-to-moderate intensity exercise after cancer treatment.³⁹ An RCT comparing high and low-to-moderate intensity endurance training within a multimodal rehabilitation program after treatment also reported no differences in CRF.⁴⁰ However, both of those studies evaluated shorter interventions after treatment, and are not directly comparable to our study.

Additional BCS did not improve CRF or other health outcomes. This may be because all groups were provided with some aspects of BCS, such as supervised training, social support, graded tasks and feedback. These methods have previously been associated with higher exercise adherence in cancer populations.^{19,20} This element of the study

design was a balance between enhancing adherence to the intervention protocol and evaluating the contribution of additional BCS. One other plausible explanation for the lack of effect of additional BCS on CRF is that participants were relatively healthy and well-motivated. It is possible that a broader, more heterogeneous clinical population would benefit from such support. Further, long-term-effects of BCS on adherence and CRF need to be examined.

Adherence to the resistance training was within range of other exercise interventions including patients undergoing curative cancer treatment,^{41,42} and exercise volume did not differ between groups. However, the adherence to the high intensity endurance training was lower than previously reported in studies with similar populations^{17,43} and lower than adherence in the low-to-moderate intensity groups. One reason for the lower levels in the present study could be that the endurance training was home-based. This decision made as we, after conducting a feasibility study and receiving input from oncology clinicians and patient representatives, deemed it would be too demanding for our participants to exercise at the gym four times per week. To facilitate adherence to the endurance training, all participants had a familiarisation period of four weeks and the average number of performed resistance and endurance sessions reached levels that could be expected to make a difference, particularly for participants with low fitness levels from start.

Another explanation for the lower adherence to high intensity endurance training compared to other studies could be that we used a stringent method for calculating adherence, including drop-outs from the intervention and taking both intensity and time into account, rather than reporting attendance only. Detailed analysis of adherence according to FITT principles in the present study has been reported in a separate publication within the research group,⁴⁴ demonstrating that once the participants had made it to the gym, they adhered to the prescribed resistance training programme to a large extent.

Post-intervention muscle strength was higher in the high intensity exercise group, which is in line with literature indicating that higher loads increase maximal strength more, even after controlling for exercise volume.⁴⁵ Although significant, the small between-group difference in cardiorespiratory fitness is consistent with the idea that not only intensity,

but also frequency, duration and volume are important exercise variables for cardiorespiratory fitness during cancer treatment.⁴⁶

Other secondary outcomes did not differ between groups. Thus, while it is well-known that exercise during cancer treatment is beneficial compared to usual care for a number of health outcomes,⁴⁷ the results of our second-generation study indicate that patients can exercise at either high or low-to-moderate intensity without missing out on improvement of several prevalent side effects.

Although only 29% of the approached patients consented to participate, the aim of this study was to compare high vs. low-to-moderate intensity exercise for CRF. As such, internal validity was prioritised over external validity (generalisability). Our sample consisted of relatively healthy and motivated individuals; mainly well-educated women treated for breast cancer. Moreover, individuals included in the main analysis had a slightly higher cardiorespiratory fitness and health related quality of life than those who dropped out after randomisation. However, baseline levels of these factors did not differ between randomisation groups for those included in the analysis. Participants varied in age, current exercise levels and perceived importance of exercise and it is unlikely that biological mechanisms related to exercise intensity differ in our sample compared with the broader population. However, we cannot claim that the results can be generalised to the total population of patients with curatively intended treatment for breast, prostate or colorectal cancer, due to the different characteristics in the sample and the small proportions of patients with prostate and colorectal cancer.

Strengths and limitations

A clear protocol and competent, trained coaches helped ensure that the intervention was delivered consistently across the different centres and the participants were closely monitored and given frequent face-to-face feedback regarding intensity. Objective measures were used to assess physical activity as well as maximal testing of cardiorespiratory fitness and muscle strength. Supplementary complete cases analysis showed largely similar results as the main analysis, suggesting that missing data were not differential. However, difference between the intention-to-treat and the complete case analysis for VO_2 max suggests that missing data for this analysis may be differential.

The study was not powered to draw conclusions about the effects of exercise intensity on CRF for specific diagnosis groups. However, diagnosis specific results for breast cancer and colorectal cancer were of the same magnitude and direction as the main results. These findings can be meta-analysed with other studies to inform research on diagnosis-specific effects. Blinding coaches and participants to the intervention was not feasible. However, there was not strong *a priori* information about which intensity would be better for CRF, so this is unlikely to introduce serious bias. Since we did not have full control of the home-based endurance training, bias may have been introduced if there was differential reporting between groups. However, this was limited by providing participants with a heart rate monitor to objectively measure intensity and duration of home-based training, rather than relying on self-report. Furthermore, adherence to the endurance training differed between groups indicating that we did not have full control of exercise volume for this intervention component.

Conclusions

For CRF, we found no clinically important difference between participants randomised to high vs low-to-moderate intensity exercise. Patients undergoing (neo-)adjuvant treatment for breast, prostate or colorectal cancer can therefore be advised to exercise at either intensity, according to their own preferences. There were few and minor adverse events during the intervention, indicating that exercise is safe, even at high intensity, for these patient groups. In a motivated and relatively healthy sample, additional BCS is not likely to influence CRF. Future studies are needed to evaluate such support in broader clinical populations.

PERSPECTIVE

It is well established that exercise during and after cancer treatment improves CRF⁷ and international guidelines recommend cancer survivors to stay active and perform endurance and/or resistance training 2-3 times per week.¹³ However, the importance of exercise intensity for CRF is unclear. The present study found that participants undergoing (neo-)adjuvant cancer treatment exercising at high intensity for 6 months demonstrated lower physical fatigue at post-intervention, compared to participants

exercising at low-to-moderate intensity. However, the difference between groups was below the threshold for clinical importance. There was no effect of behaviour change support (goal-setting, planning and self-monitoring of exercise) on fatigue. The 6-month intervention, including resistance and endurance training, caused few minor adverse events, indicating that exercise is safe even at high intensity and can be recommended to these patient groups. The important message to clinicians is that patients undergoing (neo-)adjuvant treatment for breast, prostate or colorectal cancer can be advised that it is safe to exercise at either high or low-to-moderate intensity, according to their own preferences. Behaviour change support in terms of goal-setting, planning and self-monitoring of exercise may be unnecessary if patients are relatively healthy, motivated for exercise and participate in supervised, well-controlled interventions.

Contributors

All authors were involved in drafting the article or revising it critically for important intellectual content and agreed to be accountable for all aspects of the work ensuring that questions related to the accuracy and integrity of the work were appropriately resolved.

All authors approved of the final version to be submitted.

Study conception and design: Demmelmaier, Henriksson, Mazzoni, Igelström, Ax, Sjövall, Hellbom, Pingel, Lindman, S Johansson, Velikova, Raastad, Buffart, Åsenlöf, Aaronson, Glimelius, Nygren, B Johansson, Börjeson, Berntsen, Nordin.

Acquisition of data: Demmelmaier, Henriksson, Mazzoni, Helgesen Björke, Igelström, Ax, Sjövall, Lindman, S Johansson, B Johansson, Börjeson, Berntsen, Nordin.

Analysis and interpretation of data: Demmelmaier, Brooke, Henriksson, Mazzoni, Helgesen Björke, Pingel, Velikova, Raastad, Buffart, Åsenlöf, Aaronson, Glimelius, Nygren, B Johansson, Börjeson, Berntsen, Nordin.

Funding

This work was supported by The Swedish Cancer Society (grant numbers 150841, 160483); The Swedish Research Council (grant number KDB/9514); The Nordic Cancer Union (2015), and The Oncology Department Foundations Research Fund in Uppsala (2016, 2017).

Ethical approval

The Swedish Ethical Review Authority approved the study (Dnr 2014/249). All participants gave written informed consent.

Data sharing statement

Deidentified participant data (including data dictionaries) will be shared upon reasonable request for research purposes by contacting the corresponding author.

ORCID iD

Ingrid Demmelmaier 0000-0002-2068-4708

REFERENCES

1. Arnold M, Rutherford MJ, Bardot A, et al. Progress in cancer survival, mortality, and incidence in seven high-income countries 1995-2014 (ICBP SURVMARK-2): a population-based study. *Lancet Oncol*. 2019;20(11):1493-1505.
2. Stein KD, Syrjala KL, Andrykowski MA. Physical and psychological long-term and late effects of cancer. *Cancer*. 2008;112(S11):2577-2592.
3. Bower JE. Cancer-related fatigue: Mechanisms, risk factors, and treatments. *Nature Rev Clin Oncol*. 2014;11(10):597-609.
4. Jones JM, Olson K, Catton P, et al. Cancer-related fatigue and associated disability in post-treatment cancer survivors. *J Cancer Surviv*. 2016;10(1):51-61.
5. Bjerkeset E, Röhrl K, Schou-Bredal I. Symptom cluster of pain, fatigue, and psychological distress in breast cancer survivors: prevalence and characteristics. *Breast Cancer Res Treat*. 2020;180(1):63-71.
6. Avis NE, Levine B, Marshall SA, Ip EH. Longitudinal examination of symptom profiles among breast cancer survivors. *J Pain Symptom Manage*. 2017;53(4):703-710.
7. Mustian KM, Alfano CM, Heckler C, et al. Comparison of pharmaceutical, psychological, and exercise treatments for cancer-related fatigue: a meta-analysis. *JAMA Oncol*. 2017;3(7):961-968.
8. Hojman P, Gehl J, Christensen JF, Pedersen BK. Molecular mechanisms linking exercise to cancer prevention and treatment. *Cell metabolism*. 2018;27(1):10-21.
9. Buffart LM, Kalter J, Sweegers MG, et al. Effects and moderators of exercise on quality of life and physical function in patients with cancer: An individual patient data meta-analysis of 34 RCTs. *Cancer Treat Rev*. 2017;52:91-104.
10. van Waart H, Stuiver MM, van Harten WH, et al. Effect of low-intensity physical activity and moderate- to high-intensity physical exercise during adjuvant chemotherapy on physical fitness, fatigue, and chemotherapy completion rates: results of the PACES randomized clinical trial. *J Clin Oncol*. 2015;33(17):1918-1927.

- Accepted Article
11. Courneya KS, Segal RJ, Mackey JR, et al. Effects of aerobic and resistance exercise in breast cancer patients receiving adjuvant chemotherapy: a multicenter randomized controlled trial. *J Clin Oncol*. 2007;25(28):4396-4404.
 12. McTiernan A, Friedenreich CM, Katzmarzyk PT, et al. Physical activity in cancer prevention and survival: a systematic review. *Med Sci Sports Exerc*. 2019;51(6):1252-1261.
 13. Campbell KL, Winters-Stone KM, Wiskemann J, et al. Exercise guidelines for cancer survivors: consensus statement from international multidisciplinary roundtable. *Med Sci Sports Exerc*. 2019;51(11):2375-2390.
 14. Wong JN, McAuley E, Trinh L. Physical activity programming and counseling preferences among cancer survivors: a systematic review. *Int J Behav Nutr Phys Act*. 2018;15(1):48-48.
 15. Wang Z, McLoone P, Morrison DS. Diet, exercise, obesity, smoking and alcohol consumption in cancer survivors and the general population: a comparative study of 16 282 individuals. *Br J Cancer*. 2015;112(3):572-575.
 16. Winters-Stone KM, Dobek J, Nail L, et al. Strength training stops bone loss and builds muscle in postmenopausal breast cancer survivors: a randomized, controlled trial. *Breast Cancer Res Treat*. 2011;127(2):447-456.
 17. Witlox L, Velthuis MJ, Boer JH, et al. Attendance and compliance with an exercise program during localized breast cancer treatment in a randomized controlled trial: The PACT study. *PLoS One*. 2019;14(5):e0215517.
 18. Lavallée JF, Abdin S, Faulkner J, Husted M. Barriers and facilitators to participating in physical activity for adults with breast cancer receiving adjuvant treatment: A qualitative metasynthesis. *Psycho-Oncol*. 2019;28:468-476.
 19. Bourke L, Homer K, Thaha M, et al. Interventions for promoting habitual exercise in people living with and beyond cancer. *Cochrane Database Syst Rev*. 2013;9:CD010192
 20. Turner RR SL, Quirk H, Greasley RU, Saxton JM, Taylor SJC, Rosario DJ, Thaha MA, Bourke L. Interventions for promoting habitual exercise in people living with and beyond cancer. *Cochrane Database Syst Rev*. 2018;9:CD010192.
 21. Finne E, Glausch M, Exner AK, Sauzet O, Stolzel F, Seidel N. Behavior change techniques for increasing physical activity in cancer survivors: a systematic review

- and meta-analysis of randomized controlled trials. *Cancer Manag Res.* 2018;10:5125-5143.
22. Grimmett C, Corbett T, Brunet J, et al. Systematic review and meta-analysis of maintenance of physical activity behaviour change in cancer survivors. *Int J Behav Nutr Phys Act.* 2019;16(1):37.
 23. Furmaniak AC, Menig M, Markes MH. Exercise for women receiving adjuvant therapy for breast cancer. *Cochrane Database Syst Rev.* 2016;9:CD005001.
 24. Berntsen S, Aaronson NK, Buffart L, et al. Design of a randomized controlled trial of physical training and cancer (Phys-Can) – the impact of exercise intensity on cancer related fatigue, quality of life and disease outcome. *BMC Cancer.* 2017;17(1):218.
 25. Smets EM, Garssen B, Bonke B, De Haes JC. The Multidimensional Fatigue Inventory (MFI) psychometric qualities of an instrument to assess fatigue. *J Psychosom Res.* 1995;39(3):315-325.
 26. Cella D. The Functional Assessment of Cancer Therapy-Anemia (FACT-An) Scale: a new tool for the assessment of outcomes in cancer anemia and fatigue. *Seminars in hematology.* 1997;34(3 Suppl 2):13-19.
 27. Aaronson NK, Ahmedzai S, Bergman B, et al. The European Organization for Research and Treatment of Cancer QLQ-C30: a quality-of-life instrument for use in international clinical trials in oncology. *J Natl Cancer Inst.* 1993;85(5):365-376.
 28. Zigmond AS, Snaith RP. The hospital and anxiety and depression scale. *Acta Psychiatr Scand.* 1983;67:361-370.
 29. Garin O, Ayuso-Mateos JL, Almansa J, et al. Validation of the "World Health Organization Disability Assessment Schedule, WHODAS-2" in patients with chronic diseases. *Health Qual Life Outcomes.* 2010;8:51.
 30. Longo DL, Duffey PL, DeVita VT, Jr., Wesley MN, Hubbard SM, Young RC. The calculation of actual or received dose intensity: a comparison of published methods. *J Clin Oncol.* 1991;9(11):2042-2051.
 31. Garber CE, Blissmer B, Deschenes MR, et al. American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in

- apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc.* 2011;43(7):1334-1359.
32. Edvardsen E, Hansen BH, Holme IM, Dyrstad SM, Anderssen SA. Reference values for cardiorespiratory response and fitness on the treadmill in a 20- to 85-year-old population. *Chest.* 2013;144(1):241-248.
33. Borg G. Psychophysical bases of perceived exertion. *Med Sci Sports Exerc.* 1982;14:377-381.
34. Bjørke ACH, Raastad T, Berntsen S. Criteria for the determination of maximal oxygen uptake in patients newly diagnosed with cancer: Baseline data from the randomized controlled trial of physical training and cancer (Phys-Can). *PLoS One.* 2020;15(6):e0234507.
35. Nilsen TS, Scott JM, Michalski M, et al. Novel methods for reporting of exercise dose and adherence: an exploratory analysis. *Med Sci Sports Exerc.* 2018;50(6):1134-1141.
36. Purcell A, Fleming J, Bennett S, Burmeister B, Haines T. Determining the minimal clinically important difference criteria for the Multidimensional Fatigue Inventory in a radiotherapy population. *Support Care Cancer.* 2010;18(3):307-315.
37. Enders C. Improving the accuracy of maximum likelihood analyses. In: *Applied missing data analysis.* New York, NY: The Guilford Press; 2010:127-163.
38. Enders CK. A note on the use of missing auxiliary variables in full information maximum likelihood-based structural equation models. *Struct Equ Modeling.* 2008;15(3):434-448.
39. Kampshoff CS, Chinapaw MJ, Brug J, et al. Randomized controlled trial of the effects of high intensity and low-to-moderate intensity exercise on physical fitness and fatigue in cancer survivors: results of the Resistance and Endurance exercise After ChemoTherapy (REACT) study. *BMC Med.* 2015;13:275.
40. Schmitt J, Lindner N, Reuss-Borst M, Holmberg HC, Sperlich B. A 3-week multimodal intervention involving high-intensity interval training in female cancer survivors: a randomized controlled trial. *Physiol Rep.* 2016;4(3).
41. Kirkham AA, Bonsignore A, Bland KA, et al. Exercise prescription and adherence for breast cancer: one size does not FITT all. *Med Sci Sports Exerc.* 2018;50(2):177-186.

- Accepted Article
42. Kirkham AA, Bland KA, Zucker DS, et al. "Chemotherapy-periodized" exercise to accommodate for cyclical variation in fatigue. *Med Sci Sports Exerc.* 2020;52(2):278-286.
 43. Mijwel S, Backman M, Bolam KA, et al. Adding high-intensity interval training to conventional training modalities: optimizing health-related outcomes during chemotherapy for breast cancer: the OptiTrain randomized controlled trial. *Breast Cancer Res Treat.* 2018;168(1):79-93.
 44. Mazzoni A-S, Brooke HL, Berntsen S, Nordin K, Demmelmaier I. Exercise adherence and effect of self-regulatory behavior change techniques in patients undergoing curative cancer treatment: secondary analysis from the Phys-Can randomized controlled trial. *Integr Cancer Ther.* 2020;19:1534735420946834.
 45. Kraemer WJ, Ratamess NA. Fundamentals of resistance training: progression and exercise prescription. *Med Sci Sports Exerc.* 2004;36(4):674-688.
 46. Bjørke ACH, Sweegers MG, Buffart LM, Raastad T, Nygren P, Berntsen S. Which exercise prescriptions optimize VO₂max during cancer treatment?. A systematic review and meta-analysis. 2019;29(9):1274-1287.
 47. Fairman CM, Focht BC, Lucas AR, Lustberg MB. Effects of exercise interventions during different treatments in breast cancer. *J Community Support Oncol.* 2016;14(5):200-209.

Table 1 Description of resistance training, endurance training and additional behaviour change support (BCS) components of the intervention, according to 2017 CONSORT checklist for reporting randomized trials assessing non-pharmacological treatment.

	Resistance training	Endurance training	Additional BCS
Content	<p><i>High intensity:</i> 3 x 6 RM (2 minutes rest between sets) once a week. Last set until failure. 3 x 10 RM (1 minute rest between sets) once a week. Last set until failure.</p> <p><i>Low-to-moderate intensity:</i> 3 x 12 repetitions at 50% of 6 RM (2 minutes between sets) once a week. 3 x 20 repetitions at 50% of 10 RM (1 minute rest between sets) once a week.</p> <p>Two sessions per week.</p>	<p><i>High intensity:</i> Twice-weekly interval sessions. Two minutes of exercise (running, cycling, walking up-hill) at 80-90% HRR followed by two minutes of active rest. Progression from 5 intervals, adding intervals over time until max 10 intervals. Warm-up and cool-down for 5-10 minutes respectively.</p> <p><i>Low-to-moderate intensity:</i> 150 weekly minutes of endurance activity (walking, cycling) in bouts of minimum 10 minutes at 40-50% of HRR.</p> <p>HRR was determined for each participant based on a VO₂max test performed before the intervention.</p>	<p>Coaches guided participants in using strategies to facilitate adherence to the exercise, focusing mainly on the home-based endurance training:</p> <p><i>Goal-setting</i> <i>Short-term action planning</i> <i>Self-monitoring</i> <i>Review of goal-setting</i> <i>Behavioural analysis</i> <i>Long-term coping planning</i></p>
Setting	Supervised at public gyms in groups of typically 5-10 participants. Separate groups for each of the four conditions.	Homebased.	Face-to-face concurrent with resistance training at the gym. Only <i>self-monitoring</i> was homebased and performed by the participants after endurance training. <i>Long-term coping planning</i> was performed at the end of the exercise period.

Tailoring	Individually adapted weights based on repeated testing of 6 and 10 RM in all exercises. Weights were lowered temporarily for participants struggling with severe side-effects, and then successively increased. Exercises that caused pain were substituted with other exercises activating the same muscle groups.	Individually adapted intensity based on VO ₂ max tests and heart rate monitors. Type of activity according to individual preferences, e.g. walking, bicycling, running.	Adapted to the participants' needs; fewer/short reviews if goals were easily reached every week. <i>Goal-setting</i> and <i>action planning</i> specifying when, where and how to train. Based on interviews about previous exercise habits. <i>Self-monitoring</i> by extended logbooks with facilitators and barriers in specific situations. <i>Review</i> and adjustment of <i>goal-setting</i> to be important and realistic to the participant. <i>Analysis</i> by identifying determinants of training, based on logbooks and discussions. <i>Long-term coping planning</i> according to participants' preferences about maintained physical activity/exercise.
Standardisation	Familiarisation period of 6 weeks. Six exercises; 3 for the upper extremities and three for the lower extremities. Four additional exercises for the trunk and pelvic floor were advised but not controlled (see Supplementary Figure 1). Progression based on testing of 6 and 10 RM every 4-6 weeks.	Familiarisation period of three weeks. All participants wore heart rate monitors and recorded their exertion according to the Borg RPE [27] in exercise logbooks.	Weekly reviews during the first month and then typically every 4-6 weeks, included in week-to-week checklists. Printed sheets for coaches to align the procedures for <i>goal-setting</i> , <i>action planning</i> , <i>review of goal-setting</i> , <i>analysis</i> and <i>long-term coping planning</i> . Electronic or printed extended logbooks for participants' <i>self-monitoring</i> .
Provider adherence: assessment and enhancement	Three-day course for coaches on supervising participants' exercise according to a detailed intervention protocol. Repeated on-site visits and project group	See left.	<i>BCS coaches</i> had three additional course days with theory and practice on BCS and a detailed protocol. Repeated on-site visits by research staff and project meetings on five

	meetings with research staff on five occasions. Twice-monthly tele conferences with coaches from each site to discuss and align the delivery of the intervention. Week-by-week checklist for each participant, corresponding to the intervention protocol, was used by coaches.		occasions. Audio recordings of reviews were used twice to assess the coaches' use of BCS and feedback was provided to them by research staff. <i>Non-BCS coaches</i> had a protocol specifying what they were not allowed to do. These restrictions and any problems with adhering to them were followed up repeatedly at on-site visits and project meetings.
Participant adherence: assessment and enhancement	Week-by-week checklist for each participant with attendance, 6 and 10 RM test results and notes about deviations from the protocol. Printed logbooks where target weights were recorded by the coaches. The coaches checked adherence to the protocol and gave feedback at each session. If participants did not attend a resistance training session, they were contacted by telephone and encouraged to attend the next session.	Files from heart rate monitors were reviewed by coaches together with the participants. All participants completed standardized logbooks for endurance training, either electronically or by paper. Pulse files and logbooks were checked for intensity and overall adherence and feedback was provided.	For <i>self-monitoring</i> , the extended logbooks were checked regularly by the coaches and the participants were encouraged to use them. <i>Goal-setting reviews</i> and <i>analysis</i> were performed weekly during the first month of the exercise period and then typically every 4-6 week. <i>Long-term coping plans</i> were written; one copy for the participant and one for the coach to follow up. Telephone follow-up by coach at 3 and 9 after end of the exercise period.
Calculation of adherence	Performed training divided by maximum possible training using logbook data. Performed training was performed weight x performed number of repetitions, summed across all exercises and training sessions.	Performed training divided by maximum possible training using a combination of logbook and pulse file data. <i>High intensity:</i> Performed training was number of intervals x interval duration summed across all training	Performed number of sessions including reviews and action planning divided by the maximum possible number of sessions according to the protocol (n = 9).

	<p>Maximum possible training was weight x number of repetitions according to the protocol summed across all exercises and maximum possible training sessions.</p>	<p>sessions with an average intensity of minimum 90% of the 80% lower HRR limit. This adjustment was made to take into account biking sessions; lower HR despite similar exertion level as running. Maximum possible training was number of intervals x interval duration (2 minutes) x 2 x number of weeks of training according to the protocol.</p> <p><i>Low-to-moderate intensity:</i> Performed training was minutes of activity of an intensity of 40-60% of HRR. Adjustment of upper limit was made as general heart rate increase is common during chemotherapy/cortisol treatment. Maximum possible training was 150 x number of weeks of training according to the protocol. In addition, calculation of minutes of activity at >60%, enabling adjustment for high intensity.</p>	
--	---	---	--

RM = Repetition maximum, HRR = Heart rate reserve, VO_2 max = Maximum oxygen respiratory uptake, RPE = Rating of perceived exertion

Table 2. Sociodemographic, disease and planned treatment data at baseline by intervention group for randomised participants. Data are mean (SD) or number (%). N vary due to missing data, % is of those with data available.

	HI with BCS (n=144)	HI without BCS (n=144)	LMI with BCS (n=145)	LMI without BCS (n=144)
Age, years	59.3 (13.0)	58.1 (11.4)	58.0 (11.6)	59.6 (11.8)
Sex				
Female	115 (79.9)	116 (80.6)	118 (81.4)	116 (80.6)
Living situation				
Living with partner	112 (80.0)	114 (83.2)	117 (84.2)	109 (79.6)
Education				
University	79 (56.0)	84 (60.9)	92 (65.7)	81 (58.3)
Smoking or using snuff				
Never	72 (57.6)	70 (54.3)	76 (59.8)	70 (53.4)
Previous/less than daily	44 (35.2)	53 (41.1)	47 (37.0)	47 (35.9)
Daily	9 (7.2)	6 (4.7)	4 (3.1)	14 (10.7)
Weight status				
Normal weight, BMI 18-24.9 kg/m ²	60 (45.5)	72 (52.2)	66 (49.3)	69 (50.0)
Pre-obese, BMI 25-29.9 kg/m ²	44 (33.3)	46 (33.3)	54 (40.3)	44 (31.9)
Obese, BMI>29.9 kg/m ²	28 (21.2)	20 (14.5)	14 (10.4)	25 (18.1)
Comorbidities				
Yes	79 (57.7)	78 (58.6)	77 (55.4)	93 (66.4)
Current exercise habits[†]				
Endurance training since >6 months	41 (35)	49 (40)	51 (42)	38 (30)
Resistance training since >6 months	27 (23)	25 (21)	22 (19)	14 (12)
Self-reported importance of[‡]				
HI endurance training	56 (32)	58 (34)	60 (31)	54 (34)
LMI endurance training	77 (28)	76 (27)	79 (26)	79 (22)
Resistance training	68 (29)	71 (30)	72 (29)	70 (29)
Breast cancer[§]	113	115	116	113
T in situ-T1	69 (69.7)	63 (63.6)	56 (55.4)	69 (69.7)
T2-T3	30 (30.3)	35 (35.4)	45 (44.6)	30 (30.3)
N1	15 (15.2)	16 (16.2)	15 (14.9)	17(17.2)
Chemotherapy[¶]	70 (67.3)	66 (61.1)	69 (64.5)	71 (67.0)
Adjuvant	60 (85.7)	52 (78.8)	58 (84.1)	61 (85.9)
Neoadjuvant	10 (14.3)	14 (21.2)	11 (15.9)	10 (14.1)
Antibody treatment	20 (28.6)	19 (28.8)	17 (24.6)	23 (32.4)
Radiotherapy^{††}	84 (80.8)	88 (81.5)	85 (79.4)	92 (86.8)

<i>Endocrine treatment</i>	72 (69.2)	75 (69.4)	84 (78.5)	79 (74.5)
Prostate cancer	26	23	23	25
T1-T2	20 (76.9)	19 (82.6)	17 (73.9)	18 (72.0)
T3-T4	2 (7.7)	3 (13.0)	1 (4.3)	4 (16.0)
N1	1 (3.8)	0	3 (13.0)	1 (4.0)
<i>Radiotherapy</i> ^{††}	25 (100.0)	20 (100.0)	22 (100.0)	25 (100.0)
<i>Endocrine treatment</i>	13 (50)	13 (56.5)	10 (43.5)	15 (60)
Colorectal cancer ^{§§}	5	6	6	6
T2-T4	5 (100.0)	6 (100.0)	6 (100.0)	5 (100.0)
N1-N2	5 (100.0)	4 (66.7)	4 (66.7)	5 (100.0)
<i>Chemotherapy</i> ^{¶¶}	5 (100.0)	6 (100.0)	6 (100.0)	5 (100.0)

HI: High intensity exercise; LMI: Low-to-moderate intensity exercise; BCS: Additional behaviour change support; BMI: body mass index. [†]Exercise Stage Assessment Instrument categories 1-5 with 1=Pre-contemplation stage and 5=Maintenance stage, physically active longer than 6 months. [‡]Visual analogue scale 0-100 mm anchored at 'Not at all important' and 'Very important'. T: tumour size. N: lymph node status. [§]One participant in HI without BCS had stage T4d treated with curative intent. Two participants in LMI without BCS had N2 and one in LMI with BCS had N3. [¶]Chemotherapy was Epirubicine-based and/or Taxane-based. ^{††}Breast and/or axilla. ^{‡‡}Brachy and/or external. ^{§§}One participant in HI with BCS had radically removed liver metastasis. One participant in HI with BCS and two in HI without BCS had pre-operative radiotherapy. ^{¶¶}Capecitabine-Oxaliplatin or Capecitabine only.

Table 3. Descriptive data for primary and secondary outcome measures at baseline and post-intervention (i.e. 6 month follow-up) for patients with data available

	HI with BCS		HI without BCS		LMI with BCS		LMI without BCS									
	Baseline		Follow-up		Baseline		Follow-up									
	n	mean (SD)	n	mean (SD)	n	mean (SD)	n	mean (SD)								
Primary outcome CRF																
MFI General Fatigue	134	11.1 (4.3)	107	10.2 (4.5)	135	11.3 (4.6)	105	10.2 (4.0)	139	11.6 (4.6)	110	10.4 (4.3)	136	11.4 (4.5)	117	10.8 (4.5)
MFI Physical Fatigue	136	11.3 (4.2)	112	8.8 (4.4)	134	11.0 (4.5)	107	8.3 (3.9)	139	11.1 (4.3)	106	9.5 (4.2)	138	11.3 (4.2)	111	10.2 (4.3)
MFI Reduced Activity	134	10.4 (3.9)	110	9.4 (4.4)	134	10.5 (4.2)	106	9.6 (3.9)	140	11.0 (4.1)	108	9.6 (3.8)	137	10.7 (4.1)	116	9.7 (4.1)
MFI Reduced Motivation	139	8.3 (3.5)	110	7.9 (3.4)	134	8.9 (3.4)	106	8.0 (3.2)	139	8.7 (3.6)	109	7.7 (3.2)	135	8.9 (3.3)	116	8.3 (3.6)
MFI Mental Fatigue	138	8.8 (4.1)	111	8.5 (4.2)	136	9.3 (4.0)	105	8.8 (3.8)	138	9.6 (4.2)	107	9.0 (3.9)	133	9.6 (3.9)	116	9.4 (4.2)
FACIT Fatigue subscale	140	41.9 (8.6)	112	41.8 (9.7)	137	40.8 (8.6)	108	42.5 (8.0)	139	39.9 (9.3)	112	42.2 (8.7)	141	39.6 (9.9)	116	41.9 (9.0)
Secondary outcomes																
EORTC QLQ-C30 Summary Score	135	85.2 (11.7)	111	84.4 (13.8)	136	81.8 (12.5)	106	86.6 (9.6)	133	82.5 (12.9)	111	86.6 (11.1)	134	82.4 (12.8)	112	84.9 (10.7)
HADS Depression	140	3.2 (3.2)	112	2.7 (3.2)	137	3.3 (3.2)	108	2.9 (3.2)	142	3.4 (3.2)	110	2.6 (2.8)	142	3.6 (3.2)	118	3.0 (2.9)
HADS Anxiety	140	4.9 (4.3)	112	3.9 (4.1)	137	5.9 (4.5)	108	4.0 (3.8)	142	5.6 (4.4)	110	3.9 (3.7)	142	5.8 (4.4)	118	4.0 (4.0)
WHODAS Work subscale	84	5.6 (5.1)	44	4.5 (4.7)	82	7.0 (5.8)	50	3.7 (3.4)	82	6.6 (5.4)	52	4.1 (3.7)	82	5.3 (5.1)	57	3.2 (3.9)
WHODAS Social Participation subscale	136	7.5 (5.6)	109	6.6 (5.6)	134	9.0 (5.5)	106	6.6 (4.9)	139	8.1 (5.7)	110	6.1 (5.3)	137	8.3 (5.1)	112	6.2 (4.8)
Average 1RM left and right leg, kg	118	59.7 (20.1)	93	66.7 (21.5)	119	56.2 (21.1)	97	65.5 (21.3)	122	58.8 (21.1)	99	63.0 (20.8)	119	54.9 (19.9)	109	60.3 (19.7)
1RM chest press, kg	119	32.7 (12.2)	94	36.6 (12.7)	118	32.4 (11.6)	101	36.8 (12.9)	119	34.1 (14.3)	98	37.6 (15.2)	116	31.5 (13.2)	106	34.9 (14.1)

VO ₂ max, ml/kg/min	107 30.5 (7.2)	92 30.0 (7.3)	121 30.9 (7.0)	97 30.8 (6.6)	115 31.2 (7.1)	99 30.3 (7.2)	121 29.1 (7.1)	95 28.4 (6.8)
Sleep, hrs/day	118 7.3 (1.0)	105 7.1 (0.9)	116 7.3 (0.9)	95 7.3 (1.1)	121 7.4 (1.1)	105 7.1 (1.1)	127 7.2 (1.2)	97 7.1 (1.3)
Sedentary time, hrs/day	118 18.3 (2.2)	105 18.0 (2.2)	116 18.1 (1.9)	95 17.9 (1.9)	121 18.2 (1.8)	105 18.0 (1.7)	127 18.7 (1.7)	97 18.4 (2.0)
MVPA, hrs/day	118 1.1 (0.8)	105 1.2 (0.8)	116 1.3 (1.0)	95 1.5 (1.1)	121 1.3 (0.8)	105 1.5 (0.8)	127 1.1 (0.7)	97 1.2 (0.9)
Relative Dose Intensity % [†]		75 90.2 (18.0)		72 92.6 (13.0)		75 93.6 (11.0)		76 92.9 (12.2)

[†]Relative dose intensity calculated only for patients treated with chemotherapy. HI: High intensity exercise; LMI: Low-to-moderate intensity exercise; BCS: Additional behaviour change support; SD: standard deviation; CRF: Cancer-related fatigue; MFI: Multidimensional Fatigue Inventory; FACIT: Functional Assessment of Chronic Illness Therapy; EORTC QLQ-C30: European Organisation for Research and Treatment of Cancer, Quality of life Questionnaire C30; HADS: Hospital Anxiety and Depression scale; WHODAS: World Health Organization Disability Assessment Schedule; 1RM: 1 repetition maximum; VO₂max: maximal volume of oxygen uptake; MVPA: moderate-to-vigorous intensity physical activity.

Table 4. Main effects of exercise intensity, additional behaviour change support and interaction post-intervention after multiple imputation by chained equations to account for missing data, presented as adjusted mean difference and 95% confidence intervals (n=577).

	Exercise intensity AMD (95%CI)	BCS AMD (95%CI)	Interaction AMD (95%CI)	p- value for interaction effect
Primary outcome CRF				
MFI General Fatigue	-0.36 (-1.04 to 0.33)	-0.20 (-0.91 to 0.51)	0.19 (-0.53 to 0.91)	0.608
MFI Physical Fatigue	-1.05 (-1.85 to -0.25)	-0.43 (-1.21 to 0.34)	0.26 (-0.55 to 1.08)	0.524
MFI Reduced Activity	0.22 (-0.49 to 0.92)	-0.35 (-1.04 to 0.35)	-0.05 (-0.76 to 0.67)	0.899
MFI Reduced Motivation	0.05 (-0.53 to 0.64)	-0.27 (-0.83 to 0.30)	0.31 (-0.25 to 0.87)	0.276
MFI Mental Fatigue	-0.26 (-0.94 to 0.42)	-0.20 (-0.88 to 0.48)	0.36 (-0.32 to 1.04)	0.304
FACIT Fatigue subscale	-0.43 (-1.87 to 1.01)	-0.11 (-1.59 to 1.36)	-0.63 (-2.09 to 0.84)	0.401
Secondary outcomes				
EORTC QLQ-C30 Summary Score	-0.64 (-2.42 to 1.15)	-0.77 (-2.51 to 0.97)	-2.83 (-4.61 to -1.05)	0.002
HADS Depression	0.03 (-0.45 to 0.50)	-0.24 (-0.74 to 0.27)	0.17 (-0.32 to 0.66)	0.495
HADS Anxiety	0.16 (-0.39 to 0.71)	0.23 (-0.35 to 0.81)	0.51 (-0.06 to 1.08)	0.079
WHODAS Work subscale [†]	0.31 (-1.01 to 1.63)	0.42 (-0.86 to 1.71)	0.73 (-0.59 to 2.06)	0.277
WHODAS Social Participation subscale	0.12 (-0.68 to 0.91)	0.36 (-0.45 to 1.18)	0.51 (-0.28 to 1.30)	0.206
Average 1RM left & right leg, kg [‡]	3.98 (0.58 to 7.38)	2.85 (-0.59 to 6.29)	-0.66 (-3.86 to 2.55)	0.687
1RM chest press, kg [‡]	0.41 (-1.62 to 2.44)	1.47 (-0.52 to 3.47)	-1.40 (-3.41 to 0.61)	0.171
VO ₂ max, ml/kg/min [‡]	1.61 (0.19 to 3.04)	0.76 (-0.68 to 2.20)	-1.25 (-2.67 to 0.16)	0.082
Sleep, hrs/day [‡]	0.10 (-0.15 to 0.34)	-0.03 (-0.27 to 0.22)	-0.10 (-0.33 to 0.14)	0.422

Sedentary time, hrs/day [†]	-0.26 (-0.68 to 0.16)	-0.13 (-0.54 to 0.28)	0.26 (-0.14 to 0.66)	0.204
MVPA, hrs/day [†]	0.06 (-0.14 to 0.26)	0.07 (-0.13 to 0.26)	-0.26 (-0.45 to -0.07)	0.008
Relative Dose Intensity, % [§]	-1.54 (-4.46 to 1.39)	-0.99 (-3.94 to 1.96)	-1.99 (-4.93 to 0.95)	0.184

[†]For participants who reported working. [‡]Baseline values not included in analysis due to missing data >10%. [§]For participants treated with chemotherapy. Linear regression analyses adjusted for hospital, cancer site, and baseline measure of outcome. Bold indicates p-value<0.05. AMD: Adjusted mean difference; 95%CI: 95% confidence intervals; BCS: Additional behaviour change support; CRF: Cancer-related fatigue; MFI: Multidimensional Fatigue Inventory; FACIT: Functional Assessment of Chronic Illness Therapy; EORTC QLQ-C30: European Organisation for Research and Treatment of Cancer, Quality of life Questionnaire C30; HADS: Hospital Anxiety and Depression scale; WHODAS: World Health Organization Disability Assessment Schedule; 1RM: 1 repetition maximum; VO₂max: maximal volume of oxygen uptake; MVPA: moderate-to-vigorous intensity physical activity.

Figure legends

Figure 1. CONSORT diagram of flow of participants through the Phys-Can study. Numbers with (in)complete baseline and follow-up data are based on cancer related fatigue (MFI physical fatigue subscale), exact numbers for other outcomes vary (Table 3). Follow-up refers to data collected at the post-intervention. HI: high intensity exercise, LMI: low-to-moderate intensity exercise, BCS: additional behaviour change support.

Figure 2. Adherence to prescribed strength and endurance training volume, by training group. Bars represent mean adherence, error bars indicate 1 standard deviation from the mean. Participants who dropped out of the study were recorded as 0 adherence to any remaining training sessions. P-values reflect pair-wise comparisons across the four intervention groups using Tukey post-hoc tests. All other pair-wise comparisons resulted in p-values >0.05. HI: high intensity exercise, LMI: low-to-moderate intensity exercise, BCS: additional behaviour change support.

Supplementary figure 1. Illustrations of resistance training exercises.




Figure 1. CONSORT diagram of flow of participants through the Phys-Can study. Numbers with (in)complete baseline and follow-up data are based on cancer related fatigue (MFI physical fatigue sub-scale), exact numbers for other outcomes vary (see Table 3). Follow-up refers to data collected at the end of the 6 month intervention. HI: High intensity exercise; LMI: Low-to-moderate intensity exercise, BCS; additional behaviour change support.




Figure 2. Adherence to prescribed strength and endurance training volume, by training group. Bars represent mean adherence, error bars indicate 1 standard deviation from the mean. Participants who dropped out of the study were recorded as 0 adherence to any remaining training sessions. P-values reflect pair-wise comparisons across the four intervention groups using Tukey post-hoc tests. All other pair-wise comparisons resulted in p-values >0.05. HI: high intensity exercise, LMI: low-to-moderate intensity exercise, BCS: additional behaviour change support.